

Vehicle tyres – A key source of microplastics to the environment: An introduction to the fate, effects, and mitigation strategies

This working paper was prepared independently by members of the Scientists' Coalition for an Effective Plastics Treaty: Andy M. Booth (SINTEF Ocean, Norway), Lisbet Sørensen (SINTEF Ocean, Norway), Susanne Brander (Oregon State University, US) Judith S. Weis (Rutgers University, New Jersey, US), Winnie Courtene-Jones (Bangor University), Florence Parker-Jurd (University of Plymouth), and Richard C. Thompson (University of Plymouth).

Corresponding author email: andy.Booth@sintef.no

It is estimated 12.7 MT of plastic enter the environment as microplastics every year, the main sources being paint, tyre particles, pellets, textiles and personal care products¹. There will also be substantial additional quantities from the fragmentation of larger items of plastic that have already entered the environment as debris. This document focuses on emissions from vehicle tyres which are estimated to account for an annual release of 3.6 MT of microplastic to the environment¹. Every year, almost 2 billion new vehicle tyres are produced world-wide. At the same time >1 billion tyres reach their end-of-life. During use, tyre wear particles (TWPs) are formed through abrasion of the rubber material in contact with the road surface. By mass, tyres are one of the largest sources of secondary microplastics, but reported particulate emission volumes and environmental concentrations are highly variable, both spatially and temporally. The most commonly detected size ranges of TWPs are 0.05-100 µm, with most of the particles being below 2.5 µm²⁻⁴. Tyres and the emitted TWPs are known to contain a high level and diversity of chemicals of concern, many of which have been shown to leach into environmental matrices⁵.

Source emissions: There is limited data on TWP emissions from direct measurements of environmental matrices⁶. Urban, highway and rural roads are considered the main point sources of TWP emissions, where higher emissions result from increasing traffic loads. Release per km per vehicle type typically decreases in the order lorry/18-wheeler > truck > bus > van > passenger car > motorcycle > moped (i.e., decreasing with decreasing vehicle weight)². Braking, accelerating, and turning causes increased emissions for all vehicle types⁷. Emissions per km travelled are largest in urban areas (due to increased braking, accelerating and turning), followed by highways and rural areas. It is estimated that through its usable lifetime, a tyre loses between 10-30% of its initial mass. Considering an average initial mass of 8.2 kg for a passenger car tyre and of 70 kg for a truck tyre, around 1.2 kg (passenger car) and 12.6 kg (truck) of tread weight is lost to the environment during the working lifespan of the tyre^{8,9}. However, not all tyres are fully worn before they should be replaced (ca. every 5 years). The amount of tyre wear may be higher for electric vehicles, which are heavier than those that are fuel-powered and are increasing in number⁵.

Pathways: Road run-off,^{10,11} atmospheric deposition of smaller TWPs, and re-distribution through wastewater treatment plants (WWTPs) may be considered the primary transport pathways for TWPs to reach the natural environment^{12,13}. Re-purposing of roadside soils, and repurposing of end-of-life tyres as crumb in artificial sports pitches are potentially overlooked, additional pathways of TWPs to the natural environment.

Plastic chemicals in vehicle tyres: Tyres contain a complex mixture of chemicals, including organic chemicals and heavy metals¹⁴. Both unreactive chemicals (e.g., polymers, oils, fillers, wax, resins and antioxidants) and reactive chemicals (e.g., sulphur compounds, retardants, accelerators, activators and glue) are used in the composition of the tyre tread (where TWPs originate). Literature reviews have shown that ~800 chemicals are used and/or present in tyres and TWPs¹⁵.


Chemical groups in descending order by number of known chemicals: vulcanisation accelerators, antioxidants and antiozonants, polymerisation agents, plasticiser, other protectants, adhesion agents, nitrosamine compounds, reaction products, mastication agents, blowing agents, vulcanisation agents, vulcanisation retarders, vulcanisation activators.

Notably, decomposition products and 'other' chemicals represent some of the largest groups of chemicals.

As exact tyre formulations are subject to producer's intellectual properties rights, there is limited transparency of the volumes used in different tyres.

Importantly, many additives and non-intentionally added substances (NIAS) can leach out of vehicle tyres into the surrounding environment.

However, many additive chemicals associated with tyres have other uses, making it difficult to distinguish sources based on environmental concentration measurements¹⁶⁻¹⁸. Chemical emissions from tyres can occur when the vehicle is in motion and when stationary, but still in contact with the environment (especially under wet conditions).



Overview of the emissions, transport pathways, risks and potential mitigation actions for car tyre pollutants⁵

TWP and chemical environmental transport routes and accumulation: Roadside soil is the primary recipient of TWPs^{19,20}, with an estimated 45-80% of TWP deposition occurring within the first few metres from the roadside. Small TWPs and mobile chemicals may wash-out from soils and transfer to groundwaters. Surface waters (lakes, rivers, seas) are the next largest recipient (6-26%) and can act as accumulation zones and transport mechanisms for TWPs and tyre chemicals to the marine environment when located close to main non-urban roads (e.g., highways)²¹⁻²³. Coastal towns and cities may represent major points of emission and transport of TWPs and tyre chemicals to the marine environment via urban run-off or by passing through WWTP systems. Due to their variable encrustation of road wear materials the density and therefore buoyancy of TWPs is variable^{24,25}. Furthermore, due to their small size (0.05-100 μm) their sinking rates can be quite long as drag processes become more dominant. Smaller TWPs (<100 μm) may remain airborne for longer and may be subject to long-range transport before deposition occurs⁴.

Impacts of TWPs and their plastic chemicals: Additive and NIAS chemicals leaching from tyres have been demonstrated to be a source of significant toxicity to aquatic organisms, but the complexity of the mixture makes identifying the most toxic chemicals challenging^{26,27}. For example, tyre leachates have been implicated in urban stormwater mortality syndrome in salmonids on the West coast of North America¹¹. This syndrome, which causes 40-90% of returning salmon to die before spawning, threatening salmonid conservation, was associated with degraded water quality, urbanisation, and high traffic intensity. Recently, hybrid toxicity identification evaluation and effect-directed analysis was used to identify the chemical 6PPD quinone (a degradation product of the antiozonant 6PPD) as the primary cause. The physical presence of TWPs can also be toxic to organisms, causing food dilution and additional responses similar to other microplastic types^{28,29}, but further work is needed to separate chemical and physical particle effects. In addition to aquatic systems, there is increasing evidence that TWPs may lead to changes in essential ecosystem services and multifunctionality, and therefore represent an emerging threat to soil health²⁰. Furthermore, human exposure to tyre wear and chemicals is highest near roads.

Potential Interventions:

- Simplification of tyre formulations to remove/replace the chemicals of highest concern
- Improved road runoff collection and processing
- Societal changes, including increased use of public transport, car sharing, changes in driving style, lower speed limits, reducing vehicle weight, ensuring correct tyre inflation
- Regulations on minimum tyre abrasion limits and transparent labelling to inform customer choice

Please cite this as: Booth, A., Sørensen, L., Brander, S., Weis, J.S., Courtene-Jones, W., Parker-Jurd, F., Thompson, R.C. (2024) Vehicle tyres – A key source of microplastics to the environment: An introduction to fate, effects, and mitigation strategies. Available from: <https://ikhapp.org/news-and-event/working-papers-developed-by-members-of-the-scientists-coalition/>.

References:

1. Earth Action. 2023. Leakage of microplastics into oceans and land. EA's global assessment & benchmark of literature
2. Kole, P. J. *et al.* Wear and Tear of Tyres: A Stealthy Source of Microplastics in the Environment. *International Journal of Environmental Research and Public Health* 2017, 14
3. Grigoratos, T. *et al.* Experimental investigation of tread wear and particle emission from tyres with different treadwear marking. *Atmospheric Environment* 2018, 182.
4. Jarlskog, I. *et al.* Concentrations of tire wear microplastics and other traffic-derived non-exhaust particles in the road environment. *Environment International* 2022.
5. Mayer, P. M. *et al.* Where the rubber meets the road: Emerging environmental impacts of tire wear particles and their chemical cocktails. *Science of The Total Environment* 2024, 927.
6. Mennekes, D.; Nowack, B. Tire wear particle emissions: Measurement data where are you? *Science of the Total Environment* 2022,
7. Knight, L.J. *et al.*, 2020. Tyre wear particles: an abundant yet widely unreported microplastic?. *Environmental Science and Pollution Research*, 27
8. Valentini, F.; Pegoretti, A. End-of-life options of tyres. A review. *Advanced Industrial and Engineering Polymer Research* 2022,
9. Wagner, S. *et al.* Tire wear particles in the aquatic environment - A review on generation, analysis, occurrence, fate and effects. *Water Research* 2018, 139.
10. Eisentraut, P. *et al.* Two Birds with One Stone-Fast and Simultaneous Analysis of Microplastics: Microparticles Derived from Thermoplastics and Tire Wear. *Environmental Science and Technology Letters* 2018, 5.
11. Tian, Z. *et al.* A Ubiquitous Tire Rubber-Derived Chemical Induces Acute Mortality in Coho Salmon. *Science* 2021, 371.
12. Obermaier, N.; Pistocchi, A. A Preliminary European-Scale Assessment of Microplastics in Urban Wastewater. *Frontiers in Environmental Science* 2022, 10.
13. Parker-Jurd, F.N. *et al.* 2021. Quantifying the release of tyre wear particles to the marine environment via multiple pathways. *Marine pollution bulletin*, 172
14. Andersson-Sköld, Y.; Johannesson, M.; Gustafsson, M.; Jarlskog, I.; Lithner, D.; Polukarova, M.; Strömvall, A.-M. Microplastics from tyre and road wear: a literature review; 2020.
15. Mueller, K. *et al.* Probing the chemical complexity of tires: Identification of potential tire-borne water contaminants with high-resolution mass spectrometry. *Science of The Total Environment* 2022, 802.
16. Gao, Z. *et al.* On airborne tire wear particles along roads with different traffic characteristics using passive sampling and optical microscopy, single particle SEM/EDX, and mu-ATR-FTIR analyses. *Frontiers in Environmental Science* 2022, 10.
17. Mengistu, D. *et al.* Detection and Quantification of Tire Particles in Sediments Using a Combination of Simultaneous Thermal Analysis, Fourier Transform Infra-Red, and Parallel Factor Analysis. *International Journal of Environmental Research and Public Health* 2019, 16
18. Rauert, C. *et al.* Challenges with Quantifying Tire Road Wear Particles: Recognizing the Need for Further Refinement of the ISO Technical Specification. *Environmental Science & Technology Letters* 2021, 8 (3), 231-236.
19. Sieber, R.; Kawecki, D.; Nowack, B. Dynamic probabilistic material flow analysis of rubber release from tires into the environment. *Environmental Pollution* 2020,
20. Ding, J. *et al.* Tire wear particles: An emerging threat to soil health. *Critical Reviews in Environmental Science and Technology* 2023, 53.
21. Jarlskog, I. *et al.* Traffic-related microplastic particles, metals, and organic pollutants in an urban area under reconstruction. *Science of The Total Environment* 2021, 774.
22. Muller, A. *et al.* Determination of tire wear markers in soil samples and their distribution in a roadside soil. *Chemosphere* 2022, 294.
23. Vijayan, A. *et al.* Microplastics (MPs) in urban roadside snowbanks: Quantities, size fractions and dynamics of release. *Science of The Total Environment* 2022, 851
24. Kayhanian, M. *et al.* 2012. Characteristics of road sediment fractionated particles captured from paved surfaces, surface run-off and detention basins. *Science of the Total Environment*, 439.
25. Rhodes, E.P. *et al.* 2012. Zinc leaching from tire crumb rubber. *Environmental science & technology*, 46.
26. Boisseaux, P. *et al.* 2024. Deep dive into the chronic toxicity of tyre particle mixtures and their leachates. *Journal of Hazardous Materials*, 466.
27. Sørensen, L. *et al.* 2023. Organic chemicals associated with rubber are more toxic to marine algae and bacteria than those of thermoplastics. *Journal of Hazardous Materials*, 458.
28. Cunningham, B. *et al.* Toxicity of Micro and Nano Tire Particles and Leachate for Model Freshwater Organisms. *Journal of Hazardous Materials* 2022, 429.
29. Siddiqui, S. *et al.* Internalization, Reduced Growth, and Behavioral Effects Following Exposure to Micro and Nano Tire Particles in Two Estuarine Indicator Species. *Chemosphere*, 2022 296.